

Zymotic diseases.

	Deaths, 1860.	Deaths, 1850.	Proportions, 1860.	Proportions 1850.
Cholera	985	33,074	0.28	11.87
Cholera infantum	4,804	3,960	1.35	1.45
Croup	15,188	10,706	4.25	3.84
Diarrhœa	7,847	6,366	2.20	2.28
Dysentery	10,461	20,556	2.93	7.38
Erysipelas	2,756	2,786	0.77	1.00
Fever, intermittent	4,447	964	1.25	0.35
Fever, remittent	11,102	18,496	3.11	6.63
Fever, typhoid, typhus	19,207	13,099	5.38	4.69
Fever, yellow	657	785	0.18	0.28
Influenza	387	252	0.11	0.09
Measles	3,900	2,983	1.09	1.07
Scarlatina	26,393	9,584	7.39	3.44
Smallpox	1,263	2,352	0.35	0.84
Syphilis	231	146	0.07	0.05
Thrush	554	424	0.16	0.15
Whooping-cough	8,400	5,280	2.35	1.90
Total zymotic	118,582	131,813	33.22	47.26
Other specified diseases	218,261	134,803	61.14	48.36
Violent deaths	20,115	12,174	5.64	4.36
Unknown	36,648	44,233
Grand total	393,606	323,023	100.00	100.00